

REVIEW

Using nitric oxide to treat tendinopathy

George A C Murrell

Br J Sports Med 2007;**41**:227–231. doi: 10.1136/bjism.2006.034447

Nitric oxide (NO) is a small free radical generated by a family of enzymes, the nitric oxide synthases (NOSs). Following injury to a tendon, NO is induced by all three isoforms of NOS and NOS activity is also upregulated in tendinopathy. In animal models when NOS activity is inhibited by competitive inhibitors of NOS, tendon healing is reduced. When additional NO is added, tendon healing is enhanced. In humans, in three randomised clinical trials, we have shown that NO delivered via a transdermal patch enhances the subjective and objective recovery of patients with tennis elbow, Achilles tendinosis and supraspinatus tendinosis.

induced as a result of tendon injury in this model, and/or that expression of NOS is a part of supraspinatus tendinopathy.

HUMAN ROTATOR CUFF TENDON INJURY

During surgical repair of the rotator cuff in humans, the edges of the torn tendon are excised and discarded. We have evaluated these samples and found that NOS enzyme activity was detectable in 7 of 10 human rotator cuff tendon samples. mRNA expression was demonstrated for iNOS and eNOS isoforms in each sample examined, whereas bNOS mRNA was detectable in 3 of 8 samples.¹² These results indicate that a similar phenomenon of NOS upregulation following injury occurs in humans as in rats.

Nitric oxide (NO) is a small soluble gas synthesised by a family of enzymes, the nitric oxide synthases (NOSs). Like other free radicals, in large doses NO is toxic. In smaller, physiological doses, however, it acts as a messenger molecule.

The family of NOSs consists of three isoforms regulated by a number of cofactors. eNOS (originally found in endothelial cells¹) and bNOS (originally identified in brain and neuronal tissue²) are constitutive, low-output isoforms important in blood pressure regulation and memory. Inducible NOS (iNOS) is a high-output isoform important in host defence.^{3–4}

NOS ACTIVITY FOLLOWING TENDON INJURY

There is little NOS activity in a normal, uninjured tendon. Following division, however, we found a significant (approximately fivefold) increase in NOS activity within the healing tendon in rat Achilles⁵ and supraspinatus tendons. The activity peaks at day 7 and returns to baseline at day 14. This activity was inhibitable by an NOS inhibitor (N^G-monomethyl-L-arginine). All three NOS isoforms were expressed following rat Achilles tendon division. Four days after injury, there were increases in the steady-state levels of mRNA and protein for all 3 NOS isoforms, with peaks for iNOS (23-fold increase) at days 4 and 7, eNOS (24-fold increase) at day 7 and bNOS (7-fold increase) at day 21.^{6–7}

RAT ROTATOR CUFF TENDON HEALING

We also examined NOS expression in an acute rotator cuff tendon model in the rat by creating a defect in the supraspinatus tendon with a 3 mm diameter biopsy punch. In this model, all three NOS isoforms as determined by competitive reverse transcriptase-PCR were expressed. The expression profile was slightly different from that of healing rat Achilles tendon, with bNOS expression maximum on day 4, eNOS on day 7 and iNOS on day 7.⁸

RAT ROTATOR CUFF TENDON OVERUSE MODEL

In an exercise overuse model of tendon degeneration in the rat,^{9–10} we found that iNOS, eNOS and bNOS mRNAs were overexpressed in the supraspinatus tendon of rats subjected to treadmill running at 14 days.¹¹ These results suggest that NOS activity is

WHERE DOES NO COME FROM?

In the rat Achilles tendon model, the first isoform to appear was iNOS, followed by eNOS and then bNOS. As one would expect, iNOS was expressed in macrophage-like cells and eNOS was found in endothelial cells.⁶ Interestingly, all three isoforms were expressed in fibroblast-like cells, again in a temporal fashion, with iNOS being expressed first (days 4–7), followed by eNOS (days 4–14) and bNOS (days 14–21).⁷

IS NOS EXPRESSION IMPORTANT TO TENDON HEALING?

We fed rats a competitive NOS inhibitor (N-nitro-L-arginine methyl ester) and found that rats taking this inhibitor had significantly reduced healing of their Achilles tendons compared with rats drinking its inactive enantiomer (N-nitro-D-arginine methyl ester). There was a 50% reduction in cross-sectional area of the Achilles tendon at day 7, and a corresponding 24% reduction in the failure load of the rat Achilles tendon constructs.⁵ We have also performed experiments on iNOS knockout mice and concluded that iNOS alone is not responsible for the beneficial effects of NO on tendon healing.¹³

WHAT ROLES DOES NO PLAY IN TENDON HEALING?

The experiments on animals using NOS inhibitors show that NO is important for the volume of tissue synthesised during tendon healing (fig 1). NO is likely to be important in a number of processes, including local blood flow and host defence. Work in our laboratory has identified NO to be important in collagen synthesis.¹⁴ Cultured human rotator cuff tendon cells, when exposed to exogenous NO in the form of S-nitro-N-acetyl penicillamine and when transfected with the iNOS gene via an adenovirus

Correspondence to:
Professor G A C Murrell,
Orthopaedic Research
Institute, University of New
South Wales, The St
George Hospital, Level 2, 4-
10 South Street, Kogarah,
Sydney, New South Wales
2217, Australia; admin@
ori.org.au

Accepted 14
December 2006
Published Online First 8
February 2007

Abbreviations: NO, nitric oxide; NOS, nitric oxide synthase; bNOS, brain NOS; eNOS, endothelial NOS; GTN, glyceryl trinitrate; iNOS, inducible NOS




Figure 1 Schematic diagram of the overall effects of the addition and inhibition of nitric oxide (NO) on tendon healing.

vector, incorporated more collagenase-sensitive ^3H -proline in their matrix. This increased collagen synthesis was inhibited by an NOS inhibitor.

We used microarray analysis to elucidate global gene expression after transfection with iNOS in tenocytes isolated from the injured rotator cuff tendons of humans. The expression of a wide range of genes was affected by NO, with many activated genes having known roles in healing. Of particular significance was that NOS overexpression stimulated the transcription and translation of a range of extracellular matrix genes important to the structure of connective tissues such as tendons, including collagen type I $\alpha 1$, collagen type III $\alpha 1$, collagen IV $\alpha 5$, biglycan, decorin, laminin and matrix metalloproteinase 10. These genes were also shown to respond to stimulation by the NO donor S-nitroso-N-acetyl-penicillamine in a dose-dependent manner.¹⁵ We also showed that varying levels of NO significantly affect cellular adhesion in tenocytes, a critical process during tendon repair.¹⁵

In our rat model, we delivered NO via flurbiprofen, a non-specific cyclo-oxygenase inhibitor and via NO-paracetamol. In both experiments, the addition of NO had a protective or beneficial effect on collagen organisation, tendon healing failure load and stress (load/area).^{16, 17} These rat results are consistent with cell culture findings for human tendon cells



Figure 2 Application of a glyceryl trinitrate (GTN) patch to the shoulder for supraspinatus tendinosis.

where NO enhanced collagen synthesis, and with the results from the clinical trials described below.

CLINICAL TRIALS

To determine whether additional NO might enhance tendon healing in humans, we conducted three randomised double-blind clinical trials. These trials involved the application of a

Table 1 Genes upregulated or downregulated in cultured tendon cells following exposure to nitric oxide¹⁵

Primary function	Gene	Effect on gene regulation	
Apoptosis	Immediate early response 3	Up	
	Jun D	Up	
	COP9 constitutive photomorphogenic homologue subunit 8	Up	
	Growth arrest-specific 2 like 1	Down	
Cell adhesion/communication	Protocadherin $\alpha 6$	Up	
	Catenin (cadherin-associated protein), $\delta 2$	Down	
Cell cycle control	Cell division cycle 34	Down	
Cell growth and/or maintenance	Potassium inwardly rectifying channel, subfamily J, member 8	Up	
	Translocated promoter region (to activated MET)	Up	
	Human cellular proto-oncogene (c-met)	Down	
	TGF- β inducible early protein	Down	
Extracellular matrix/regulation	S100 calcium-binding protein A11	Up	
	Matrix metalloproteinase 10 (stromelysin 2)	Down	
	Collagen type I, $\alpha 1$	Down	
	Decorin	Down	
	Laminin B2	Down	
	Biglycan	Down	
	Collagen type III, $\alpha 1$	Down	
	Collagen type IV, $\alpha 5$	Up	
	Microfibrillar-associated protein 3-like	Up	
	Oxidation resistance 1	Up	
Signal transduction	TNF receptor-associated factor 5	Up	
	Rho GTPase-activating protein 4	Up	
	Interleukin 18 receptor 1	Up	
	IGF-II mRNA-binding protein 3	Up	
	SMA D5	Up	
	Phospholipase D1, phosphatidylcholine-specific	Up	
	Tyrosine 3-mono-oxygenase/tryptophan 5-mono-oxygenase activation protein, ζ polypeptide	Up	
	Ras suppressor protein 1	Up	
	Interferon regulatory factor 3	Up	
	A kinase (PRKA) anchor protein 10	Up	
	Mitogen-activated protein kinase 7	Down	
	Transport, cell growth and/or maintenance	Solute carrier family 22	Up

COP, coat protein complex; GTPase, guanosine triphosphatase; IGF, insulin-like growth factor; MET, methionine; PRKA, protein kinase A-anchoring; SMA, smooth muscle actin; TGF, transforming growth factor; TNF, tumour necrosis factor.

Table 2 Effects of glyceryl trinitrate 1.25 mg/day through transdermal patch and rehabilitation versus rehabilitation alone on tennis elbow (adapted from Paoloni *et al*⁹)

Week	Active	Placebo	p Value
Pain with activity—all patients (n=35) (n=39)			
2	2.1	2.3	0.31
6	1.8	1.9	0.57
12	1.5	1.6	0.61
24	1.3	1.5	0.56
Pain with activity—bilateral patients excluded (n=31) (n=37)			
2	2.1	2.4	0.11
6	1.8	1.9	0.57
12	1.5	1.6	0.72
24	1.3	1.5	0.43
Tenderness—by elbow (n=39) (n=41)			
2	1.0	1.1	0.25
6	0.7	1.1	0.014
12	0.7	1.1	0.006
24	0.5	1.0	0.014
Tenderness—bilateral data averaged (n=35) (n=40)			
2	1.0	1.2	0.08
6	0.6	1.1	0.010
12	0.6	1.1	0.003
24	0.5	1.0	0.008
Tenderness—bilateral patients excluded (n=31) (n=38)			
2	0.9	1.2	0.037
6	0.8	1.3	0.008
12	0.8	1.3	0.002
24	0.6	1.1	0.008
Change from baseline in total work—by elbow (n=39) (n=41)			
2	8.9	-1.8	0.048
6	21.6	5.6	0.007
12	23.0	6.5	0.019
24	35.9	15.4	0.019
Change from baseline in total work—all patients—bilateral data averaged (n=35) (n=39)			
2	9.8	-2.0	0.036
6	23.1	6.5	0.007
12	23.2	7.3	0.027
24	38.5	17.1	0.020
Change from baseline in total work—bilateral patients excluded (n=31) (n=37)			
2	11.0	-2.4	0.028
6	25.0	7.4	0.007
12	23.4	8.0	0.044
24	41.9	18.9	0.019
Pain at night—all patients (n=35) (n=39)			
2	1.2	1.4	0.28
6	0.9	1.2	0.20
12	0.9	1.1	0.33
24	0.7	0.8	0.44
Pain at night—bilateral patients excluded (n=31) (n=37)			
2	1.1	1.4	0.13
6	0.9	1.2	0.24
12	0.9	1.1	0.38
24	0.6	0.8	0.39
Pain at rest—all patients (n=35) (n=39)			
2	1.1	1.2	0.84
6	0.8	1.0	0.32
12	0.9	1.0	0.54
24	0.9	1.0	0.66
Pain at rest—bilateral patients excluded (n=31) (n=37)			
2	1.0	1.1	0.56
6	0.8	1.1	0.27
12	0.9	1.0	0.69
24	0.9	1.0	0.67
Change from baseline in peak force—by elbow (n=39) (n=41)			
2	1.3	0.0	0.06
6	2.5	0.8	0.016
12	2.6	0.6	0.020

Table 2 Continued

Week	Active	Placebo	p Value
24	3.4	1.4	0.025
Change from baseline in peak force—all patients—bilateral data averaged (n=35) (n=39)			
2	1.4	-0.01	0.041
6	2.6	0.9	0.015
12	2.6	0.7	0.024
24	3.7	1.6	0.024
Change from baseline in peak force—bilateral patients excluded (n=31) (n=37)			
2	1.6	-0.1	0.027
6	2.8	1.0	0.016
12	2.6	0.8	0.036
24	4.1	1.8	0.021

All results are adjusted for the baseline (week 0) value.

commercially available NO delivery system (glyceryl trinitrate (GTN) patches) and a placebo version of the same patch. These patches were applied over the area of tenderness for three conditions: tennis elbow, Achilles tendinosis and supraspinatus tendinosis (fig 2). A total of 53–86 patients were randomised to two groups in each of the trials. In each trial, the active group received continuous topical NO donation (1.25 mg/24 h GTN) and the placebo group received the identical patch without GTN. The

Table 3 Effects of glyceryl trinitrate 1.25 mg/day through transdermal patch and rehabilitation versus rehabilitation alone in patients with Achilles tendonitis (adapted from Paoloni *et al*⁸)

Week	Active	Placebo	p Value
Pain with activity—all patients (n=29) (n=33)			
2	1.4	1.6	0.43
6	1.1	1.4	0.22
12	0.9	1.4	0.09
24	0.6	1.0	0.15
Pain at rest—all patients (n=28) (n=33)			
2	1.2	1.1	0.70
6	0.9	1.1	0.47
12	0.9	1.1	0.31
24	0.8	0.9	0.58
Pain at night—all patients (n=27) (n=33)			
2	1.5	1.7	0.41
6	1.0	1.2	0.27
12	0.6	0.9	0.058
24	0.6	0.7	0.57
Pain on hopping—by heel (n=37) (n=44)			
2	3.2	3.2	0.98
6	2.9	3.1	0.68
12	2.3	3.0	0.10
24	1.2	2.1	0.023
Tenderness—all patients (n=29) (n=33)			
2	1.7	1.7	0.75
6	1.4	1.3	0.68
12	0.8	1.3	0.055
24	0.7	0.7	0.99
Change from baseline in total work—by heel (n=36) (n=42)			
2	12.1	12.3	0.49
6	12.9	13.2	0.66
12	13.5	13.6	0.90
24	14.9	13.6	0.10
Change from baseline in peak force—by heel (n=36) (n=41)			
2	2.2	2.2	0.41
6	2.4	2.4	0.54
12	2.6	2.4	0.30
24	2.9	2.5	0.010

All results are adjusted for the baseline (week 0) value.

Table 4 Effects of glyceryl trinitrate 1.25 mg/day through transdermal patch and rehabilitation versus rehabilitation alone in patients with supraspinatus tendinopathy (adapted from Paoloni *et al.*²⁰)

Week	Active	Placebo	p Value
Pain with activity—by shoulder			
	(n=28)	(n=28)	
2	2.2	2.1	0.80
6	1.9	2.1	0.34
12	1.6	1.9	0.36
24	1.0	1.8	0.006
Pain at rest—by shoulder			
	(n=27)	(n=28)	
2	1.7	1.9	0.47
6	1.5	1.6	0.68
12	1.1	1.4	0.19
24	0.7	1.2	0.09
Pain at night—by shoulder			
	(n=27)	(n=27)	
2	1.4	1.6	0.61
6	1.4	1.5	0.72
12	1.0	1.5	0.09
24	0.8	1.4	0.031
Tenderness—by shoulder			
	(n=27)	(n=28)	
2	0.1	0.4	0.012
6	0.2	0.5	0.017
12	0.2	0.4	0.18
24	0.2	0.2	0.60
Change from baseline in abduction range of motion—by shoulder			
	(n=27)	(n=29)	
2	7.7	1.5	0.15
6	7.8	2.0	0.29
12	16.3	1.2	0.020
24	26.6	8.7	0.004
Change from baseline in adduction power—by shoulder			
	(n=27)	(n=28)	
2	5.6	1.3	0.24
6	7.7	-0.6	0.08
12	11.9	0.4	0.023
24	9.9	1.6	0.11
Change from baseline in external rotation power—by shoulder			
	(n=27)	(n=28)	
2	4.7	0.5	0.17
6	7.4	0.8	0.11
12	11.5	0.5	0.023
24	13.4	0.3	0.013
Change from baseline in external rotation range of motion—by shoulder			
	(n=27)	(n=27)	
2	1.9	6.7	0.13
6	4.7	1.9	0.45
12	7.7	3.3	0.26
24	13.2	8.5	0.17
Change from baseline in forward flexion—by shoulder			
	(n=36)	(n=41)	
2	7.8	0.8	0.11
6	6.5	-2.3	0.09
12	13.4	-2.2	0.019
24	20.3	6.3	0.027
Impingement external rotation (% with 0)—by shoulder			
	(n=27)	(n=28)	
2	40.7	32.1	0.51
6	40.7	32.1	0.57
12	70.4	57.1	0.36
24	70.4	50.0	0.15
Impingement internal rotation (% with 0)—by shoulder			
	(n=27)	(n=28)	
2	33.3	10.7	0.059
6	48.2	17.9	0.040
12	51.9	32.1	0.15
24	74.1	39.3	0.018
Change from baseline in internal rotation power—by shoulder			
	(n=27)	(n=28)	
2	4.9	-0.9	0.14
6	10.8	2.0	0.055
12	11.0	-1.5	0.029
24	13.9	1.1	0.018
Change from baseline in internal rotation range of motion—by shoulder			
	(n=27)	(n=28)	
2	-2.3	0.5	0.37
6	-4.3	-1.0	0.22
12	-5.0	0.6	0.15
24	-8.1	-0.3	0.32

Table 4 Continued

Week	Active	Placebo	p Value
Change from baseline in subscapularis power—by shoulder			
	(n=27)	(n=28)	
2	9.0	0.9	0.009
6	9.2	5.4	0.24
12	14.0	5.8	0.045
24	15.7	6.6	0.018
Change from baseline in supraspinatus power—by shoulder			
	(n=27)	(n=28)	
2	3.6	-1.5	0.10
6	11.0	0.7	0.009
12	15.4	2.0	0.010
24	16.2	2.9	0.007

GTN patches were applied to the area of maximal tenderness once a day. Both the patients and the clinical examiner were blinded as to which group the patients were in. In each trial, the patients also received education and exercises—that is, the GTN treatment was on top of “best practice”. The results of these trials can be found in Paoloni *et al.*^{18–20}

Dr Eugene Heyman completed an independent reanalysis of the data from the three studies. In this analysis any missing data, whenever it occurred, were filled in by carrying forward the last value. Each analysis was adjusted for the baseline value. For categorical data with responses of 0–4, the baseline value was treated as a block—that is, all the patients with a baseline value of 0 were grouped together, those with a baseline of 1 were grouped together and so on. The mean results were then compared between treatment groups within each baseline score and then




Figure 3 Supraspinatus tendinosis. Effects of glyceryl trinitrate (GTN, n=28) 1.25 mg/d via transdermal patch versus rehabilitation alone (placebo, n=29) on shoulder pain with activities (A) and on dynamometer-measured supraspinatus force (B); a between-groups comparison of means and the standard error of the mean. Significant differences: *p<0.05, **p<0.01, ***p<0.001 compared with baseline using student’s paired *t* tests. (Adapted from Paoloni *et al.*²⁰).

What this study adds

- This study, involving three clinical trials, showed that delivering nitric oxide via a patch enhances clinical recovery of tendinopathy in humans.
- The enhancement is manifested by a reduction in pain, an increase in range of motion and an increase in strength.

What is already known on this topic

- Nitric oxide (NO) is important to tendon healing.
- Nitric oxide synthase (NOS) activity is upregulated in tendinopathy.
- In animal models where NOS activity is inhibited, tendon healing is reduced.
- When additional NO is added, tendon healing is enhanced.

the treatment differences were averaged while using the total number of patients within the baseline category to weight the results. Patients within the same baseline score were treated the same, but the treatment differences were weighted by the total number of patients within that baseline score. For continuous data (eg, total work), the change from baseline was used and then included as a covariate to account for any baseline differences and to account for the relationship between the baseline and the change from baseline. When patients with bilateral data were identified, the data were analysed in three ways. First, the results from the two sides were averaged, then, all bilateral patients were excluded. Finally, each group was treated separately. Tables 1–4 presents the results.

TENNIS ELBOW

The NO group had less tenderness and could perform more work and had greater peak power on the Orthopaedic Research Institute-Ankle Strength Testing System testing (table 2). The changes were most apparent at week 24. In all, 81% of patients receiving GTN patches were asymptomatic in activities of daily living at 6 months compared with 60% of patients with tendon rehabilitation alone ($p = 0.005$ with χ^2 analysis).¹⁹

ACHILLES TENDONITIS

The NO (GTN) group performed significantly better on hop testing and could generate more peak force at week 24 (table 3). In all, 78% of patients receiving GTN patches were asymptomatic for activities of daily living at 6 months compared with 49% of patients with tendon rehabilitation alone ($p = 0.001$ with χ^2 analysis). The mean effect size for all outcome measures was 14%.¹⁸

SUPRASPINATUS TENDINOPATHY

This trial produced the most significant effects. The NO group had significantly reduced shoulder pain with activity and at night, improved range of motion in abduction, forward flexion and external rotation, and improved power in abduction, external rotation, subscapularis and supraspinatus (table 4).²⁰ The changes in supraspinatus power were the most dramatic, and were significant at 6 weeks (table 4, fig 3). In all, 46% of patients receiving GTN patches were asymptomatic for activities of daily living at 6 months compared with 24% of patients with tendon rehabilitation alone ($p = 0.007$). The mean effect size of GTN treatment for all outcome measures was 26%.²⁰

DISCUSSION

NO is important to tendon healing. All three isoforms of NOS, the enzyme that produces NO, are expressed by fibroblasts

during tendon healing. Our data in animal studies, cell culture and clinical trials support the hypothesis that NO enhances extracellular matrix synthesis and results in injured tendons having better material and mechanical properties—that is, the healing tendons are stronger on a per unit area basis than those not exposed to additional NO. The clinical trials show that delivering NO via a patch enhances the clinical recovery of tendinopathy, which is manifested by a reduction in pain, an increase in range of motion and an increase in strength.

ACKNOWLEDGEMENTS

I thank the following for their invaluable contributions: Tom Best, Allison Collins, Ed Lilly, Anthony Seaber, Richard Goldner, Kim Hayes, Judie Walton, Zoltan Szomor, Jian-Hao Lin, Min Wang, Ai Wei, Wei Zhu, Ashish Diwan, Danny Jang, Xian-Hao Deng, Jo Hannafin, Helen Davies, Marty Dolan, Dedee Murrell, Richard Appleyard, Janice Nelson, Fiona Bonar, Gong-Yao Tang, Jun Yuan, Piero Del Soldato, Wei Zhu, Wei Xia, Yao Wang, Denis King, James Urbaniak, Martin Francis, Robert Duthie, Ed Puzas, Charles Turner, Chris Evans, Bruce Caterson, Csaba Szabo, Laura Manfield, Justin Paoloni, Tim Molloy, Eugene Heyman, Robert Ang, National Institutes of Health Grant AR42729 (GACM), the Soft Tissue Research Fund, Piedmont Foundation, Orthopaedic Research and Education Foundation (OREF), St George Private Hospital/Health Care of Australia and St George Hospital/South East Health, and NiCOX Corporation and Cure Therapeutics.

Competing interests: None declared.

REFERENCES

- 1 Palmer RMJ, Ashton DS, Moncada S. Vascular endothelial cells synthesize nitric oxide from L-arginine. *Nature* 1988;**333**:664–6.
- 2 Bredt DS, Hwang PM, Snyder SH. Localization of nitric oxide synthase indicating a neural role for nitric oxide. *Nature* 1990;**347**:768–70.
- 3 Stuehr DJ, Nathan CF. Nitric oxide: a macrophage product responsible for cytostasis and respiratory inhibition in tumor target cells. *J Exp Med* 1989;**169**:1543–55.
- 4 Xie QW, Cho HJ, Calaycay J, et al. Cloning and characterization of inducible nitric oxide synthase from mouse macrophages. *Science* 1992;**256**:225–8.
- 5 Murrell GAC, Szabo C, Hannafin JA, et al. Modulation of tendon healing by nitric oxide. *Inflamm Res* 1997;**46**:19–27.
- 6 Lin J-H, Wang M-X, Wei A, et al. Temporal expression of nitric oxide synthase isoforms in healing Achilles tendon. *J Orthop Res* 2001;**19**:136–42.
- 7 Lin J-H, Wang M-X, Wei A, et al. The cell specific temporal expression of nitric oxide synthase isoforms during Achilles tendon healing. *Inflamm Res* 2001;**50**:1–8.
- 8 Szomor ZL. *Nitric oxide in rotator cuff tendon* [PhD thesis]. Sydney: University of New South Wales, 2003.
- 9 Carpenter JE, Flanagan CL, Thomopoulos S, et al. The effects of overuse combined with intrinsic or extrinsic alterations in an animal model of rotator cuff tendinosis. *Am J Sports Med* 1998;**26**:801–7.
- 10 Carpenter JE, Thomopoulos S, Flanagan CL, et al. Rotator cuff defect healing: a biomechanical and histologic analysis in an animal model. *J Shoulder Elbow Surg* 1998;**7**:599–605.
- 11 Szomor Z, Appleyard RA, Murrell GAC. Overexpression of nitric oxide synthases in tendon overuse. *49th Annual Meeting of the Orthopaedic Research Society*. New Orleans, LA, 2003:146.
- 12 Szomor ZL, Wang MX, Kruller A, et al. Differential expression of cytokines and nitric oxide synthase expression in human rotator cuff bursae. *Ann Rheum Dis* 2001;**60**:431–2.
- 13 Xia W, Wang Y, Appleyard R, et al. Achilles tendon healing in iNOS knockout mice. *49th Annual Meeting of the Orthopaedic Research Society*, New Orleans, LA, 2003:824.
- 14 Xia W, Szomor Z, Wang Y, et al. Nitric oxide enhances collagen synthesis in cultured human tendon cells. *J Orthop Res* 2006;**24**:159–72.
- 15 Molloy TJ, de Bock C, Wang Y, et al. Gene expression changes in SNAP-stimulated and iNOS-transfected tenocytes—expression of extracellular matrix genes and its implications for tendon healing. *J Orthop Res* 2006;**24**:1869–82.
- 16 Tang GY, Murrell GAC, Yuan J, et al. Addition of NO via NO-paracetamol enhances collagen content and organization of healing rat Achilles tendon. *Proceedings of the 50th Annual Meeting of the Orthopaedic Research Society*, New Orleans, LA, 2004:876.
- 17 Yuan J, Murrell GAC, Wei AQ, et al. Addition of nitric oxide via nitrofuribiprofen enhances the material properties of healing Achilles tendon. *Inflamm Res* 2003;**52**:230–7.
- 18 Paoloni J, Appleyard R, Murrell GAC. A randomized double-blind placebo controlled clinical trial investigating the use of topical nitric oxide application in the treatment of Achilles tendonitis. *J Bone and Joint Surg (Am)* 2004;**86-A**:916–22.
- 19 Paoloni JA, Appleyard RC, Nelson J, et al. Topical nitric oxide application in the treatment of chronic extensor tendinosis at the elbow: a randomized, double-blinded, placebo-controlled clinical trial. *Am J Sports Med* 2003;**31**:915–20.
- 20 Paoloni JA, Appleyard RC, Nelson J, et al. Topical glyceryl trinitrate application in the treatment of chronic supraspinatus tendinopathy: a randomized, double-blinded, placebo-controlled clinical trial. *Am J Sports Med* 2005;**33**:806–13.